1

4

2

5

3

ENZIMAS DE ESCAPE

ENZIMA	NOMBRE
ALT	Aspartato Amino Transferasa
TGP	Transaminasa Glutámico Pirúvica
FAS (4)	Fosfatasa Alcalina Sérica (Isoenzimas)
LDH(5)	Lactato Deshidrogenasa (Isoenzimas)
GGT	Gama Glutamil Transferasa
SD	Sorbitol Deshidrogenasa

6

1. Glutamic pyruvic transaminase (= Alanine aminotransferase)	GPT ALT	2.6.1.2	+++
2. Glutamic oxaloacetic transaminase (= Aspartate aminotransferase)	AST GOT	2.6.1.1	+
3. Glutamate dehydrogenase	GDH	1.4.1.3	++
4. Gamma-glutamyl transferase	γ -GT	2.3.2.2	++
5. Lactate dehydrogenase	LDH	1.1.1.27	+
6. Cholinesterase	ChE	3.1.1.8	+++
7. Phosphohexoisomerase	PHI	5.3.1.9	+
1. Alkaline phosphatase	AP LAP	3.1.3.1	+++
2. Leucine arylamidase (= Leucine aminopeptidase)	LAP	3.4.11.1	++
3. Gamma glutamyl transferase	γ -GT	2.3.2.2	++
4. 5'-Nucleotidase	5'-NU	3.1.3.5	+-

Tab. 5.4: Seven essential hepatocellular and four biliary diagnostic enzymes together with their respective terminology, abbreviations, EC code and specificity (+ $\rightarrow$ +++)

7

Loss of enzyme activity	4°C	20-25°C
Alkaline phosphatase	0% (10%)	
Cholinesterase	0% (0%)	
GDH	5% (15%)	
GOT	8% (12%)	
GPT	10% (17%)	
γ -GT	0% (0%)	
LAP	0% (0%)	
LDH	8% (2%)	

Stability of substrates	4°C	20-25°C
Ammonia (in hours)	2	(-)
Autoimmune antibodies	14	(5)
Cholesterol	6	(6)
Copper	14	(14)
Electrophoresis	6	(6)
Hepatitis serology	14	(5)
Immunoglobulins	7	(3)
Iron	7	(4)
Neutral fats	3	(-)
Potassium	14	(14)
Protein	6	(6)
Sodium	14	(14)

Indicator enzymes	GPT, GOT, GDH, LDH
Secretory enzymes	ChE
Excretory enzymes	AP, LAP, γ -GT, 5'-NU

Unilocular	GPT, LDH
Cytoplasm	GDH
Mitochondria	
Endoplasmic reticulum	γ -GT, ChE

Bilocular	cGOT (20%)
Cytoplasm	mGOT (80%)
Mitochondria	

Periportal (zone 1)	GPT, LDH, AP
Pericentral (zone 3)	GDH, ADH
Zones 1-3	GOT

half-life	in hours	half-life	in days
LDH ₁	10 $\pm$ 2	γ -GT	3-4
GOT	13 $\pm$ 1	AP	3-7
GDH	16 $\pm$ 2	ChE	<10
GPT	47 $\pm$ 10	LAP	<10

Tab. 5.5: Schematic differentiation of hepatocellular enzymes according to their type and preferred localization within the liver cell or the liver lobule as well as their half-life

10

Indicator enzymes	GPT, GOT, GDH, LDH
Secretory enzymes <td>ChE</td>	ChE
Excretory enzymes <td>AP, LAP, γ-GT, 5'-NU</td>	AP, LAP, γ -GT, 5'-NU

Unilocular	GPT, LDH
Cytoplasm	GDH
Mitochondria	
Endoplasmic reticulum	γ -GT, ChE

Bilocular	cGOT (20%)
Cytoplasm	mGOT (80%)
Mitochondria	

Periportal (zone 1)	GPT, LDH, AP
Pericentral (zone 3)	GDH, ADH
Zones 1-3	GOT

half-life	in hours	half-life	in days
LDH ₁	10 $\pm$ 2	γ -GT	3-4
GOT	13 $\pm$ 1	AP	3-7
GDH	16 $\pm$ 2	ChE	<10
GPT	47 $\pm$ 10	LAP	<10

Tab. 5.5: Schematic differentiation of hepatocellular enzymes according to their type and preferred localization within the liver cell or the liver lobule as well as their half-life

8

TRANSAMINASA GLUTAMICO PIRUVICA (TGP)
ALANINA AMINO TRANSFERASA (ALT)

- 1000 veces serica
- Vida media 24 h,
- magnitud directamente proporcional al daño
- Niveles permanecen por 12 horas
- Significado 2-3 veces de aumento
- Hasta 20-40 veces

ALT₁

ALT₂

- Musculo esquelético
- Igual LDH

11

Table 12.2. Cellular or tissue sources and half-lives of common serum enzymes

Enzyme	Major mechanisms that lead to increased serum activity	Major cellular sources of increased serum enzyme activity	Half-lives*
FAS	Induction	¹ Hepatocytes (L-ALP) (C-ALP in dogs) ² Biliary epithelium (I-ALP) Osteoblasts (B-ALP)	= 3 days (canine L-ALP) ^{1,2} < 8 hr (feline L-ALP) ^{1,2}
ALT	Cell damage	¹ Hepatocytes ² Skeletal myocytes	2-3 days (canine) ^{1,2}
AMS	Cell damage Decreased renal clearance	¹ Pancreatic acinar cells	= 5 hr (canine) ^{1,2}
AST	Cell damage	¹ Hepatocytes ² Skeletal myocytes Cardiac myocytes Erythrocytes	7-8 days (equine) ^{1,2,3,6} < 1 day (canine) ^{1,2,7,8}
CK	Cell damage	¹ Skeletal myocytes Cardiac myocytes Smooth muscle myocytes (intestinal)	= 2 hr (equine) ^{1,2,9} < 2 hr (canine) ^{1,2}
GGT	Increased production	¹ Biliary epithelial cells ² Hepatocytes	= 3 days (equine) ¹⁰
ID	Cell damage	¹ Hepatocytes	= 4 hr (canine) ¹¹
LD	Cell damage	¹ Hepatocytes ² Skeletal myocytes	= 6 hr (canine) ¹¹
LDH	Cell damage	¹ Skeletal myocytes Cardiac myocytes Erythrocytes	
LPS	Cell damage Decreased renal clearance	¹ Pancreatic acinar cells Liver neoplasms	= 2 hr (canine) ¹²

* Data from available but limited sources.
(¹⁻¹²) = numbers of sources in reference list.

9

FOSFATASA ALCALINA SÉRICA (FAS)

- ISOENZIMAS:
- CANINOS 4: v= 2-3 días
- Hueso (B-ALP), hepática (L-ALP),
- inducida por corticosteroides (C-ALP)
- Origen desconocido

• FELINO 2: v=8 h

• hueso, hígado

12

HEPATOPATÍA INDUCIDA POR CORTICOSTEROIDES

ESTEROIDES
Altos niveles

↓

Inducción enzimática
DNA—RNA—Proteína

↓

Daño a hepatocitos
Canaliculos

637 Histologic appearance of canine glucocorticoid hepatopathy. Note the vacuolated cells that contained glycogen prior to fixation.

13

R= ALT/FAS = 1.89

R<2 → COLESTASIS
R=2-5 Mixto colestasis y hepatocelular
R>5 HEPATOCELULAR

CANINO
AST (6-70)
ALT (4-73)
FAS (65-300)
GGT (1-9.7)

ELEVACIÓN
ALT /70 = 1.65
FAS / 389 = 0.87

Arriba del límite superior
2-3 VECES LEVE
3-4 MODERADO
>5 SEVERO

ALT/73 = 2.73
FAS/300 = 1.33

FAS.- Las isoenzimas más comunes que se encuentran en el suero de los perros son: la isoenzima hepática, la inducida por corticosteroides y la ósea.

16

GAMA GLUTAMIL TRANSFERASA (GGT)

- Derivada del hígado (también riñón y páncreas)
- Felinos, mejor que FAS
- Lipoidosis hepática felina FAS +++, GGT+
- En general aumentan de 2-3 veces significativo
- Principalmente con fibrosis porque aumenta la colestasis por el desorden de la fibrosis

- ALTAMENTE ESPECÍFICA PARA COLESTASIS (JUNTO CON FAS)
- >90% Aumento de FAS y GGT
- 50% FAS
- 80% GGT

14

GGT, unida a membrana celular: tubulo proximal, hígado, páncreas, intestino, bazo
HIGADO: principalmente cél. Conductos biliares
MUY SENSIBLE DE DAÑO A CONDUCTOS, poco específica
Muy útil para diferenciar origen hepático de la FAS (no hueso, no esteroides)

FIBROSIS HEPÁTICA

17

Woreta & Alqahtani

Fig. 2. Concentration of AST and ALT according to location of hepatocytes in portal triad.

Si ALT normal
AST/ALT =>5
daño extrahepático

VIDA MEDIA
AST= 12-22 horas
ALT = 37-57 h

RECUPERACIÓN:
ALT>AST

CPK
LDH

CANINO
AST (6-70)
ALT (4-73)
FAS (65-300)
GGT (1-9.7)

AST/ALT = 0.65

15

Fig. 4. Algorithm for evaluation of patients with elevated alkaline phosphatase. ERCP, endoscopic retrograde cholangiopancreatography.

18

► Generally, the elevation of transaminases is a reliable marker of hepatocellular damage. (19–21, 23, 24, 28, 32, 34)

GOT/GPT ratio (or DeRitis ratio)

- Inflammatory type: GOT < GPT < 1
- Necrotizing type: GOT > GPT > 1
- In severe alcoholic or malignant liver disease: GOT > GPT > 2

γ-GT/GPT ratio

- Acute viral hepatitis: < 1
- Toxic hepatitis: < 2
- Chronic persistent hepatitis: < 2–3
- Chronic hepatitis: Acute alcoholic hepatitis Cirrhosis: < 3–6
- Alcoholic cirrhosis: < 3–6
- Acute obstructive jaundice: > 6
- Biliary cirrhosis: > 6
- Chronic obstructive jaundice: > 6
- Liver carcinoma: > 12
- Liver metastases: > 12

GPT + GOT/GDH ratio

- Severe toxic/inflammatory damage of peritonitis/pneumonia: < 10
- Hepatic metastases: < 10
- Obstructive jaundice: < 20
- Necrotic episode in acute hepatitis: Acute intoxication, liver carcinoma: < 30–50
- Biliary cirrhosis: < 30–40
- Acute episode in chronic hepatitis or in liver cirrhosis: < 30–50
- Cholestasis: < 30–50
- Acute viral hepatitis: < 30
- Acute alcoholic hepatitis: < 30

Tab 5.6: Enzyme ratios (using optimized tests) as additional "cheap" and "fast" enzyme parameters of differential diagnosis of liver diseases (13, 18, 33)

LDH/GOT ratio

- Haemolytic jaundice: > 12
- Hepatocellular jaundice: < 12

GPT/GDH ratio

- Obstructive jaundice: < 10
- Hepatocellular jaundice: > 10

Tab 5.7: Enzyme ratios as additional support for the differential diagnosis of hyperbilirubinemia

► **Elevated GPT is the most sensitive indicator of hepatocellular damage. An exclusive rise in GPT is, as a rule, of hepatocellular origin.**

Tab 5.8: Diagnostic accuracy in the detection or exclusion of hepatopathy using 7 hepatobiliary enzymes in 520 patients suffering from various liver diseases (15)

1. GPT (= screening enzyme)	= 81%
2. GPT + γ-GT + CHE	= 95%
3. GPT + γ-GT + CHE + AP (= 3-fold pattern)	= 97%
4. GPT + γ-GT + CHE + AP + GOT + GDH (= 5-fold pattern)	= 97%
5. GPT + γ-GT + CHE + AP + GOT + GDH + LDH (= 7-fold pattern)	= 97%

19

Box 1

Primary hepatobiliary tumor types and examples

- Hepatocellular
 - Hepatic adenoma
 - Hepatocellular carcinoma
 - Hepatoblastoma
- Biliary
 - Biliary adenoma (or cystadenoma)
 - Biliary carcinoma
- Neuroendocrine
 - Neuroendocrine carcinoma or carcinoid
- Mesenchymal
 - Hemangiomasarcoma
 - Leiomyosarcoma
 - Fibrosarcoma
 - Osteosarcoma
 - Malignant mesenchymoma
 - Chondrosarcoma

22

Tabla 2: Hepatosidad por drogas.

I. Daño hepatocelular:	II. Colestasis:
<ul style="list-style-type: none"> Acetaminofén (Tylenol). Azathioprina (Imuran). Allopurinol (Zyloric). Diclofenaco (Voltaren). Hydralazina. Isoniazida. Methotrexate. Methylopa (Aldomet). Nitrofurantoina (Furadantin). Quinidina. Estatinas. Corticosteroides. Anidoclorona (Cardaron). 	<ul style="list-style-type: none"> Esteroides anabólicos. Amoxicilina-clavulanato (Augmentin). Captopril (Capoten). Clozapina. Entromonina. Estrogenos. Contraceptivos orales. Fenitina. Carbamazepina. Sulfas. Tetraciclina. Acido Valproico.

20

23

Enzimas Hepáticas

UNIVERSIDAD DE GUAYAQUIL
Facultad de Ciencias Químicas
Carrera de Química y Farmacia
Asignatura: Análisis Químico Clínico II

TRANSAMINASAS

Aspartato-aminotransferasa

Enzima ampliamente utilizada en el estudio de las enfermedades hepáticas. No obstante, se encuentra presente en células de diversos tejidos, tales como el músculo esquelético y cardíaco, riñón, eritrocitos y leucocitos, entre otros. Por que hacer el análisis? Para el cribado y diagnóstico de enfermedad hepática.

Alanino-aminotransferasa

La prueba de ALT se suele realizar como una parte del estudio general de la función hepática, que también incluye el análisis de otras enzimas hepáticas. La determinación de la concentración de ALT junto con la de otras enzimas hepáticas como la aspartato-aminotransferasa, o AST, y la fosfatasa alcalina, o ALP, proporcionan a los médicos una información más específica sobre los problemas hepáticos.

FOSFATASA ALCALINA

La fosfatasa alcalina es una enzima que cataliza la hidrólisis del enlace éster fosfórico en un grupo orgánico y un grupo fosfórico a pH alcalino, lo que libera fosfato al medio. Está presente en riñón, hígado, intestino y hueso. La medida de niveles anormales de fosfatasa alcalina en el suero indican la existencia de enfermedades más digestivas (traumatismo, osteomielitis, hiperparatiroidismo secundario, osteosarcoma) o haca datos hepáticos.

GAMMA GLUTAMIL TRANSPEPTIDASA

Es una enzima de membrana, siendo el marcador más sensible de enfermedad hepática, proviene fundamentalmente del hígado y del tejido sano. Algunas alteraciones como la enfermedad de Pájar, cirrosis, neoplasias vesicales, hiperparatiroidismo y ictericos pueden elevar ligeramente los niveles de fosfatasa alcalina en suero.

Valores Normales

- Mujeres: Hasta 31 U/L
- Hombres: Hasta 40 U/L

Valores Normales

- Mujeres: Hasta 31 U/L
- Hombres: Hasta 37 U/L

Valores Normales

- 98 a 279 U/L

Valores Normales

- 7 - 50 U/L

Nombre: Rodríguez Panchana Roberto
G-I
Fecha: 11/08/2021

21

24

25

Box 2
Potential extrahepatic causes for elevated serum or plasma activities of alanine aminotransferase and aspartate aminotransferase

- Diabetes mellitus
- Enteritis
- Hemolysis (AST)
- Hyperthyroidism (cats)
- Hypoxia: cardiac disease, hepatic thrombosis, anemia
- Hypothyroidism (dogs)
- Myopathy (more likely AST)
- Pancreatitis/pancreatic neoplasia
- Peritonitis
- Septicemia
- Vacuolar hepatopathy (secondary to exogenous/endogenous glucocorticoids)

28

γ -glutamyltransferase
 GGT is an enzyme that catalyzes the transfer of the γ -glutamyl group of peptides, such as glutathione to other peptides or amino acids. GGT is present in the cell membranes of many tissues including the proximal renal tubule, liver, pancreas, intestine, and spleen.¹ In the liver, GGT is located primarily on biliary epithelial cells and on the apical membrane of hepatocytes. The predominant source of serum GGT is the liver. Entry of GGT into the serum may occur by solubilization and release of membrane-bound GGT or the death of biliary epithelial cells.²

Serum GGT is a sensitive indicator of the presence of injury to the bile ducts or liver. However, its use is limited by its lack of specificity, because many nonhepatic disorders can lead to elevation, including diabetes, hyperthyroidism, chronic obstructive pulmonary disease, and renal failure.³ Alcohol abuse and certain medications, such as barbiturates or phenytoin, lead to induction of hepatic microsomal GGT.^{1,4} The main clinical use of GGT is to confirm the hepatic origin of elevated AP levels, because GGT is not elevated in patients with bone disease.

26

World Small Animal Veterinary Association (WSAVA)

ACUMULACIÓN DE GLUCÓGENO

Fig. 10. Corticosteroid hepatopathy in a dog. Excessive endogenous or exogenous corticosteroids can produce distended hepatocytes with irregular clear vacuoles and diaphanous strands of cytoplasm. The vacuoles contain glycogen that is removed during fixation and processing.

29

Box 1
Potential hepatobiliary causes for elevated serum or plasma activities of alanine aminotransferase and aspartate aminotransferase

- Drug-induced liver injury: for example, acetaminophen, anesthetic agents, arsenical compounds, carprofen, diazepam/oxazepam, griseofulvin, itraconazole, ketoconazole, lomustine, phenobarbital, phenytoin, primidone, mebendazole, methimazole, oxibendazole-diethylcarbamazine, tetracycline, doxycycline, clindamycin, nitrofurantoin, trimethoprim-sulfadiazine, azathioprine
- Hepatic lipidosis (cats)
- Infectious: for example, ascending enteric bacterial infection, feline infectious peritonitis, schistosomiasis, leptospirosis
- Inflammatory: copper-associated chronic hepatitis, idiopathic chronic hepatitis, cholangitis, gall bladder mucocele
- Lysosomal storage disease
- Neoplasia (primary or metastatic)
- Nodular hyperplasia
- Portosystemic shunt (congenital or acquired)
- Toxin ingestion: for example, aflatoxin, amanita mushroom, blue-green algae, copper, herbicides, insecticides, iron, sago palm, zinc, xylitol
- Trauma
- Vacuolar hepatopathy (idiopathic)

27

Fig. 11. Macrovesicular steatosis in a dog. Lipid vacuoles are distinct, round, and clear. Typically, these vacuoles displace the nucleus to the edge of the cell.

30

Fig. 12. Microvesicular steatosis in a cat. Microvesicular lipid vacuoles are smaller than the nucleus but retain a clear round outline, and the nucleus remains central in affected cells.

31

34

Fig. 13. Hepatic amyloidosis in a cat. Uniformly hyaline eosinophilic material within the sinusoids distends the sinusoids and leads to atrophy of the hepatocytes that remain.

32

SÍNTESIS

- Proteínas (15% diferentes) [la mitad]
- Factores de coagulación (vit)
- Colesterol
- Ac Grasos
- Lipoproteínas
- Urea
- Factor eritropoyético hepático

35

FUNCIONES DEL HÍGADO

- Producción de sustancias a la sangre.

33

36

SÍNTESIS

- Proteínas (15%)
- Factores de coagulación
- Colesterol
- Ac Grasos
- Lipoproteínas
- Urea
- Factor eritropoyético

37

Electroforesis de Proteínas Plasmáticas

40

PRUEBAS DE LABORATORIO

Proteínas

Tabla 15. Valores del análisis electroforético del suero de vacas con afecciones hepáticas (según VENTUROLI, 1958)

Enfermedad	Albumina (%)	Globulina (%)			Proteína total (g/100 ml)	Cociente albúmina/globulina
		α	β	γ		
Animales sanos	43.5	14.5	13.5	27.0	6.4	0.77
Parásitos hepáticos	34.7	14.8	15.8	35.3	6.1	0.52
Equinocosis hepática	33.7	15.7	15.0	35.8	6.2	0.51
Afección por cuerpo extraño	29.9	13.6	18.7	37.8	6.5	0.42
Tuberculosis hepática	22.9	14.2	17.7	45.2	8.3	0.29

38

ELECTROFORESIS DE PROTEÍNAS

Figure 4—Serum protein electrophoresis from a normal, healthy cat.

Figure 5—Serum protein electrophoresis from a cat with nonrefractive FIP. Hypergammaglobulinemia is evident.

41

Determinación de Proteínas

- Proteína total (BHC)
- Proteína total, albúmina, globulinas relación A/G
- Electroforesis de proteínas séricas.
- Proteínas en orina (tira reactiva)
- Proteínas en orina (químico, electroforesis)

39

SIGNOS CLÍNICOS

42

43

46

44

47

45

Table 30-4

LIPID AND LIPOPROTEIN CHANGES IN LIVER DISEASE

Liver Disease	Lipid and Lipoprotein Changes
Parenchymal Disease	
Acute	<ul style="list-style-type: none"> ↑ Triglycerides ↓ Triglycerides* ↓ Cholesterol esters ↓ VLDL, ↓ HDL ↓ LCAT, ↓ HL, ± ↓ LPL
Chronic	<ul style="list-style-type: none"> ↑ free: ester cholesterol* ↓ VLDL, ↓ HDL, ↓ LCAT, ↓ HL
Major Bile Duct Occlusion	<ul style="list-style-type: none"> ↑ Cholesterol (free and total)* ↓ Cholesterol esters* ↓ LCAT, ↑ Lipoprotein X*
Portosystemic Shunting	<ul style="list-style-type: none"> ↓ Cholesterol (total)* ↓ Triglycerides*

* Humans and dogs

VLDL: Very low density lipoproteins

HDL: High density lipoproteins

LCAT: Lecithin cholesterol acyl transferase

HL: Hepatic lipase

LPL: Lipoprotein lipase

48

344. A cat with liver disease displays numerous spur projections on the erythrocytes (acanthocytosis) (arrows).

Trigliceridos
Colesterol
Acidos Grasos
Urea

Otras causas

FALTA EN DIETA
INSUFICIENCIA EN ABSORCIÓN
DESORDENES METABÓLICOS (DIABETES)

49

52

FUNCIONES DEL HÍGADO

- Eliminación de sustancias de la sangre (destoxificación).

50

53

EXCRECIÓN

- Bilirrubinas
- Hormonas, Esteroides
- Fármacos
- Anestésicos
- Antibióticos
- Toxinas
- Complejos amoniacales NH_4^+

51

PRUEBAS DE LABORATORIO

- Bilirrubinas : tot, dir, ind. (ictericia >2-3 mg/dl)
- BSF.- 1g, 5 y 12 min. Vida media, (2.5-4.1)
- <5% 30 min.
- Amoniaco.
- Sales biliares

54

Bilirrubinas

55

FUNCIONES DEL HÍGADO

- Digestión (Ac.o sales biliares, bicarbonato ¿FAS?)

58

Bilirrubinas

56

FUNCIONES DE ACIDOS BILIARES

- Digestión de grasas
- Absorción de vitaminas liposolubles
- Homeostasis del colesterol
- Se originan del colesterol y glicerol
- Taurocólico, queno-deoxicólico
- Se reabsorben en intestino 95% (hasta 15 veces al día)

59

SIGNOS CLÍNICOS

>1.6-1.8 mg/dl
 NORMAL Bil t <1.1 mg/dl (<18.8 umol/L)
 Bil i (0.3- 1.0 mg/dl (5-17 umol/L))

57

60

DIGESTIÓN

- Sales Biliares
- (Fosfatasa alcalina)
- Bicarbonato

61

PRUEBAS DE LABORATORIO

- Examen Coprológico
- Turbidez del plasma
- Ac. biliares suero
- Perros > 30mM/L (<10)
- Gatos > 20 mM/l (<5)

64

62

FUNCIONES DEL HÍGADO

- Almacén de sustancias.

65

DIGESTIÓN

- Sales Biliares
- (Fosfatasa alcalina)
- Bicarbonato

63

66

ALMACÉN

- FISIOLÓGICO
 - Glucógeno
 - Vitaminas
 - Hierro
 - Selenio
 - Cobre
- PATOLÓGICO
 - Grasa
 - Glicoproteínas
 - Agua
 - Hierro
 - Pigmentos
 - Toxinas

67

70

68

71

69

LIPOIDOSIS

- Biosíntesis en hígado
- Rumiantes NO a partir de glucosa
- Formación alta de TG
- No exportación de TG

Fosfolípidos, triglicéridos, colesterol
Lípidos de Muy baja densidad.

72

CAUSAS DE LIPOIDOSIS

- Alto consumo de AG
- Alta producción de tej. Adiposo por demanda (Pico de lactación, finales de gestación, anorexia, Cetosis, endocrinopatías)
- Insuficiencia hepática en oxidación de AG
- Baja síntesis de apoproteínas y no exporta VLDL
- Falta de secreción de VLDL por hepatotoxinas o drogas
- Aumento de glucosa e insulina que produce altos TG, almacenados
- Alto ingreso de glucosa con alta producción de TG
- Hígado graso bovino: obesos, pocos días antes del parto, con anorexia

73

INFLAMACIÓN

Proteínas de inflamación Temprana
Elimina patógenos En sangre de origen Intestinal (Kupffer)

EXCEPTO BOVINOS
Intravascular, alvéolos pulmonares

76

FUNCIONES DEL HÍGADO

- "Inflamatoria"

74

PROTEÍNAS DE FASE AGUDA

- Más sensibles que leucocitosis o neutrofilia.
- Muchas inflamaciones las aumentan
- Canino: PCR, Haptoglobina, ceruloplasmina

77

INFLAMMATION

Anti-inflammatory Cytokines
IL-1 IL-6 TNF

Positive Acute Phase Reactants
Fibrinogen
Serum amyloid A
C-reactive protein

Haptoglobina
C3

Liver

75

78

79

ALTERACIÓN EN LAS PRUEBAS DE LABORATORIO

FASE	LESIÓN	FUNCIÓN	AGENTE
I	++		+
II	+++++	++++	++
IV	++	++	-
III	+	(+++++)	- /+++

82

80

Propietario: Sobranos
 Paciente: Molly
 Especial: Fisiol
 Edad: 13 Años
 Sexo: Hembra
 Hora: 18:24:33
 Fecha: 8 abr 2024
 Diagn: A Queen Corresponde

RESULTADO	VALORES DE REFERENCIA
Perfil Completo	
Glicemia	7.98 mmol/L ↓↓
Urea	10.29 mmol/L ↓↓
Creatinina	110 µmol/L ↓↓
Colectanol	3.22 mmol/L ↓
Bilirrubina total	14.01 µmol/L ↑↑↑
Bilirrubina conjugada	11.91 µmol/L ↑↑↑
Bilirrubina no conjugada	2.10 µmol/L ↑↑↑
AST	2381 U/L ↑↑↑
AST	624 U/L ↑↑
Fosfatasa alcalina	2324 U/L ↑↑↑
Triglicéridos	9.77 mmol/L ↑↑
CK	238 U/L ↑
Proteínas totales	56 g/L ↓
Albumina	29 g/L ↓
Globulina	35 g/L ↓
Ratión albumina / globulina	0.83 calculado
Calcio	2.75 mmol/L ↓
Fósforo inorgánico	1.56 mmol/L ↓
Potasio	4.05 mmol/L ↓
Sodio	158.0 mmol/L ↓
Cloro	124.0 mmol/L ↓
Bicarbonato	17 mmol/L ↓
Ácidos Orgánicos	39 calculado
Amilasa	161 U/L ↓
Ratión creatinina / urea	0.69 calculado
Ratión Bc/Bcr	5.67 calculado
Ratión Ca / P	2.27 calculado
Diferencia de iones fuertes	31 calculado

NO HEPÁTICO

LESIÓN hepatocelular y colestasis

ICTERICIA
 PREHEP *bil i <1
 HEP bili+bilc 1
 POST *bil c >1

83

CLASIFICACION DE LAS PRUEBAS DE LABORATORIO POR SU USO CLINICO

LESION	FUNCION	AGENTE ETIOLOGICO
Histología	Histoquímica	Inmunohistoquímica
Citología	Hormonas	Anticuerpos
Enzimas de Escape	Pruebas de Funcionamiento	Cultivos
Radiografía simple	Radiografía con medio de contraste	PCR
Examen microscópico	Examen microscópico	Hibridación
Ultrasonido	Química sanguínea	Western Blotting
		ELISA, IP, IF

81

RESULTADO	VALORES DE REFERENCIA
Hemograma	
Hematocrito	0.31 L/L
Hemoglobina	107 g/L
Eritrocitos	5.8 x10 ¹² /L
Volumen Globular Medio	46 fL
CGMH	346 g/L
Plaquetas	318 x10 ⁹ /L
Proteínas	66 g/L
Reticulocitos	0 x10 ⁹ /L
Diferencial	
Leucocitos Totales	8.20 x10 ⁹ /L
Neutrófilos Totales	6.15 x10 ⁹ /L
Neutrófilos en Banda	0.00 x10 ⁹ /L
Metametrófilos	0.00 x10 ⁹ /L
Linfocitos	0.00 x10 ⁹ /L
Monocitos	0.08 x10 ⁹ /L
Eosinófilos	0.57 x10 ⁹ /L
Basófilos	0.00 x10 ⁹ /L
E. Nucleados	0 /100 Leucocitos
Neutrófilos Tóxicos	Negativo
Linfocitos Atípicos	Negativo
Morfología de Eritrocitos	Normal
Hemoparásitos	Negativo

84